

The Lattice Λ Parameter in Domain Wall QCD

Sinya Aoki^a and Yoshinobu Kuramashi^b

^a*Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan*

^b*Institute of Particle and Nuclear Studies,*

High Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305-0801, Japan

(Dated: February 7, 2008)

Abstract

We evaluate the ratio of the scale parameter Λ in domain wall QCD to the one in the continuum theory at one loop level incorporating the effect of massless quarks. We show that the Pauli-Villars regulator is required to subtract the unphysical massive fermion modes which emerge in the fermion loop contributions to the gluon self energy. Detailed results are presented as a function of the domain wall height M .

I. INTRODUCTION

The domain-wall fermion formalism provides us an opportunity to realize the chiral symmetry at finite lattice spacing with any number of quark flavors[1, 2]. Since it is well known that the chiral symmetry plays an essential role in low energy QCD, this point is a superior feature over the Wilson and the Kogut-Susskind quark actions to perform lattice QCD simulations. While in the past two decade both conventional quark actions have been widely used to make unquenched QCD simulations incorporating the dynamical quark effects, recent progress in computational technologies allows us to embark on a full QCD simulation with the domain-wall fermion[3].

One primary step in the full QCD simulation is to determine the renormalized coupling constant, which is a fundamental parameter in QCD. At the one-loop level the renormalized coupling in the continuum $\overline{\text{MS}}$ scheme $g_{\overline{\text{MS}}}(\mu)$ at the scale μ and that on the lattice g_0 are related by

$$\frac{1}{g_{\overline{\text{MS}}}^2(\mu)} = \frac{1}{g_0^2} + \left(d_g + \frac{22}{16\pi^2} \log(\mu a) \right) + N_f \left(d_f - \frac{4}{48\pi^2} \log(\mu a) \right) \quad (1)$$

where d_g and d_f are renormalization constants: the former depends on the gauge action and the latter on the quark action. These constant terms determine the ratio of the lattice Λ parameter and the continuum one at the one-loop level. From the expressions of the Λ parameters,

$$\Lambda_L = \frac{1}{a} (\beta_0 g_0^2)^{-\beta_1/2\beta_0^2} \exp \left(-\frac{1}{2\beta_0 g_0^2} \right), \quad (2)$$

$$\Lambda_{\overline{\text{MS}}} = \mu (\beta_0 g_{\overline{\text{MS}}}^2(\mu))^{-\beta_1/2\beta_0^2} \exp \left(-\frac{1}{2\beta_0 g_{\overline{\text{MS}}}^2(\mu)} \right) \quad (3)$$

with

$$\beta_0 = \frac{1}{16\pi^2} \left(11 - \frac{2}{3} N_f \right), \quad (4)$$

$$\beta_1 = \frac{1}{(16\pi^2)^2} \left(102 - \frac{38}{3} N_f \right), \quad (5)$$

we obtain the ratio

$$\frac{\Lambda_{\overline{\text{MS}}}}{\Lambda_L} = (a\mu) \exp \left(-\frac{1}{2\beta_0} \left\{ \frac{1}{g_{\overline{\text{MS}}}^2(\mu)} - \frac{1}{g_0^2} \right\} \right) + O(g_0^2), \quad (6)$$

$$= \exp \left(-\frac{1}{2\beta_0} \{ d_g + N_f d_f \} \right) + O(g_0^2). \quad (7)$$

Thus the finite renormalization between the coupling constants defined by different regularization schemes is to relate their Λ parameters. In Table I we list the renormalization constant d_g , which has been evaluated for various gauge actions[4, 5, 6, 7, 8].

In this paper we make a perturbative determination of d_f for the domain-wall fermion, which is a necessary ingredient to extract the renormalized coupling constant from the numerical simulation with domain-wall QCD. While the perturbative techniques are well developed in our previous publications[9, 10, 11, 12, 13, 14], a characteristic feature in this calculation is to require the Pauli-Villars field to subtract the unphysical massive fermion modes, which emerge from the fermion loop diagrams.

This paper is organized as follows. In Sec. II we introduce the domain-wall fermion action with the Pauli-Villars regulator. We give their Feynman rules relevant for the present calculation. In Sec. III we evaluate the quark loop contributions to the gluon self energy in the continuum and on the lattice. The ratio of continuum and lattice Λ parameters is given in Sec. IV. Our conclusions are summarized in Sec. V.

The physical quantities are expressed in lattice units and the lattice spacing a is suppressed unless necessary. We take $SU(N_c)$ gauge group with the gauge coupling constant g .

II. ACTION AND FEYNMAN RULES

We explain the domain-wall fermion action accompanied with its Pauli-Villars regulator in the present calculation. For the fermion action we take Shamir's form[1]:

$$\begin{aligned}
S_{\text{DW}} = & \sum_n \sum_{s=1}^{N_s} \left[\frac{1}{2} \sum_{\mu} \left(\bar{\psi}(n)_s (-r + \gamma_{\mu}) U_{\mu}(n) \psi(n + \hat{\mu})_s + \bar{\psi}(n)_s (-r - \gamma_{\mu}) U_{\mu}^{\dagger}(n - \hat{\mu}) \psi(n - \hat{\mu})_s \right) \right. \\
& + \frac{1}{2} \left(\bar{\psi}(n)_s (1 + \gamma_5) \psi(n)_{s+1} + \bar{\psi}(n)_s (1 - \gamma_5) \psi(n)_{s-1} \right) + (M - 1 + 4r) \bar{\psi}(n)_s \psi(n)_s \left. \right] \\
& + m \sum_n \left(\bar{\psi}(n)_{N_s} P_R \psi(n)_1 + \bar{\psi}(n)_1 P_L \psi(n)_{N_s} \right), \tag{8}
\end{aligned}$$

where U_{μ} is the link variable of the $SU(N_c)$ gauge group. We label a four-dimensional space-time coordinate by n and an extra fifth-dimensional one by s , which runs from 1 to N_s . Since no gauge interaction is imposed along the fifth dimension, we are also allowed to consider that the index s labels the N_s “flavor” space. The parameter m denotes the physical quark mass and at $m = 0$ one chiral zero mode is realized under the condition $0 < M < 2$ for the

Dirac “mass” (domain-wall height) M . The Wilson parameter is set to $r = -1$. $P_{R/L}$ are projection operators defined by $P_{R/L} = (1 \pm \gamma_5)/2$.

The “physical” quark fields on the four-dimensional space-time are constructed in terms of the fermion fields at the boundaries,

$$q(n) = P_R \psi(n)_1 + P_L \psi(n)_{N_s}, \quad (9)$$

$$\bar{q}(n) = \bar{\psi}(n)_{N_s} P_R + \bar{\psi}(n)_1 P_L, \quad (10)$$

with which we can extract the zero mode of domain-wall fermion at $m = 0$.

We also introduce the Pauli-Villars field to subtract the massive modes of the domain-wall fermion, which emerge from the quark loop diagram in proportion to N_s in the large N_s limit. The action is given by

$$S_{PV} = S_{DW}(\bar{\psi} \rightarrow \bar{\phi}, \psi \rightarrow \phi, m = -1), \quad (11)$$

where ϕ is the N_s -flavor Wilson-Dirac boson, satisfying the anti-periodic boundary condition in the fifth direction because of the special value $m = -1$ [15]. Here we have to remark one point: no Pauli-Villars field has been introduced in the previous calculations for the renormalization factors of the composite operators[9, 10, 11, 12, 13, 14], since it requires no quark loop diagram.

For the gauge action we employ the following form:

$$S_{\text{gluon}} = \frac{1}{g^2} \left\{ c_0 \sum_{\text{plaquette}} \text{Tr} U_{pl} + c_1 \sum_{\text{rectangle}} \text{Tr} U_{rtg} + c_2 \sum_{\text{chair}} \text{Tr} U_{chr} + c_3 \sum_{\text{parallelogram}} \text{Tr} U_{plg} \right\}, \quad (12)$$

where the first term represents the standard plaquette action, and the remaining terms are six-link loops formed by a 1×2 rectangle, a bent 1×2 rectangle (chair) and a 3-dimensional parallelogram. The coefficients $c_0, \dots, c_3$ satisfy the normalization condition

$$c_0 + 8c_1 + 16c_2 + 8c_3 = 1. \quad (13)$$

There are several choices for these parameters: $c_1 = c_2 = c_3 = 0$ (Plaquette), $c_1 = -1/12$, $c_2 = c_3 = 0$ (Symanzik)[5, 16] $c_1 = -0.331$, $c_2 = c_3 = 0$ (Iwasaki), $c_1 = -0.27$, $c_2 + c_3 = -0.04$ (Iwasaki') [17], $c_1 = -0.252$, $c_2 + c_3 = -0.17$ (Wilson)[18] and $c_1 = -1.40686$, $c_2 = c_3 = 0$ (DBW2)[19]. The last four cases are called the RG-improved gauge action, whose parameters are chosen to be the values suggested by some approximate renormalization

group analyses. Some of these actions are now getting widely used to simulate the domain-wall QCD[20, 21], since they are expected to realize smooth gauge field fluctuations which approximate the continuum gauge field better than the naive plaquette action.

To prepare the Feynman rule we develop weak-coupling perturbation theory by writing the link variable in terms of the gauge potential

$$U_\mu(n) = \exp \left(i g a T^A A_\mu^A \left(n + \frac{1}{2} \hat{\mu} \right) \right), \quad (14)$$

where T^A ($A = 1, \dots, N_c^2 - 1$) is a generator of color $SU(N_c)$. Here we omit the expression of the gluon propagator and vertices, since they are not relevant for the present calculation. Quark-gluon vertices are identical to those in the N_s flavor Wilson fermion. We need one- and two-gluon vertices for the present calculation:

$$V_{1\mu}^A(q, p)_{st} = -g T^A \{ i \gamma_\mu \cos(q_\mu/2 + p_\mu/2) + r \sin(q_\mu/2 + p_\mu/2) \} \delta_{st}, \quad (15)$$

$$= -g T^A v_{1\mu}(q, p) \delta_{st}, \quad (16)$$

$$V_{2\mu\nu}^{AB}(q, p)_{st} = \frac{a}{2} g^2 \frac{1}{2} \{ T^A, T^B \} \{ i \gamma_\mu \sin(q_\mu/2 + p_\mu/2) - r \cos(q_\mu/2 + p_\mu/2) \} \delta_{\mu\nu} \delta_{st}, \quad (17)$$

$$= \frac{a}{2} g^2 \frac{1}{2} \{ T^A, T^B \} v_{2\mu\nu}(q, p) \delta_{st}, \quad (18)$$

where p and q represent incoming and outgoing quark momentum, respectively.

The fermion propagator is obtained by inverting the domain-wall Dirac operator. In the present one-loop calculation we will take $N_s \rightarrow \infty$ to neglect the contribution of the $O(e^{-\alpha N_s})$ terms with the positive constant α given in eq.(26). To this end it is sufficient to consider the following component of the fermion propagator:

$$S_F(p)_{st} \equiv \langle \psi(-p) \bar{\psi}(p) \rangle_{st} \quad (19)$$

$$= \left(-i \gamma_\mu \sin p_\mu + W^- \right)_{su} G_R(u, t) P_+ \\ + \left(-i \gamma_\mu \sin p_\mu + W^+ \right)_{su} G_L(u, t) P_-, \quad (20)$$

where the sum over the same index is taken implicitly. $W^{+/-}$ and $G_{R/L}$ are given by

$$W^+(p)_{st} = \begin{pmatrix} -W(p) & 1 & & \\ & -W(p) & \ddots & \\ & & \ddots & 1 \\ m & & & -W(p) \end{pmatrix}, \quad (21)$$

$$W^-(p)_{st} = \begin{pmatrix} -W(p) & & & m \\ & 1 & -W(p) & \\ & & \ddots & \ddots \\ & & & 1 & -W(p) \end{pmatrix}, \quad (22)$$

$$W(p) = 1 - M - r \sum_{\mu} (1 - \cos p_{\mu}). \quad (23)$$

$$G_R(s, t) = \frac{A}{F} \left[-(1 - m^2) (1 - W e^{-\alpha}) e^{\alpha(-2N_s + s + t)} - (1 - m^2) (1 - W e^{\alpha}) e^{-\alpha(s + t)} \right. \\ \left. - 2W \sinh(\alpha) m (e^{\alpha(-N_s + s - t)} + e^{\alpha(-N_s - s + t)}) \right] + A e^{-\alpha|s - t|}, \quad (24)$$

$$G_L(s, t) = \frac{A}{F} \left[-(1 - m^2) (1 - W e^{\alpha}) e^{\alpha(-2N_s + s + t - 2)} - (1 - m^2) (1 - W e^{-\alpha}) e^{\alpha(-s - t + 2)} \right. \\ \left. - 2W \sinh(\alpha) m (e^{\alpha(-N_s + s - t)} + e^{\alpha(-N_s - s + t)}) \right] + A e^{-\alpha|s - t|}, \quad (25)$$

$$\cosh(\alpha) = \frac{1 + W^2 + \sum_{\mu} \sin^2 p_{\mu}}{2|W|}, \quad (26)$$

$$A = \frac{1}{2W \sinh(\alpha)}, \quad (27)$$

$$F = 1 - e^{\alpha} W - m^2 (1 - W e^{-\alpha}). \quad (28)$$

In case that W becomes negative the fermion propagator is given with the replacement $e^{\pm\alpha} \rightarrow -e^{\pm\alpha}$. In the present calculation we consider only $m = 0$ case. The Feynman rule for the Pauli-Villars regulator is obtained by setting the quark mass to $m = -1$.

III. FERMION LOOP CONTRIBUTION TO THE GLUON SELF-ENERGY

We first present the calculation of the vacuum polarization diagram in Fig. 1 (a) employing a continuum regularization scheme with the massless quarks. Here we choose the naive dimensional regularization (NDR) as the ultraviolet regularization scheme. Both the Dirac matrices and the fermion loop momenta of the Feynman integrals are defined in $D = 4 - 2\epsilon$ ($\epsilon > 0$) in this scheme. To regularize the infrared divergence we take the external gluon momentum p to be off-shell $p^2 \neq 0$.

The gluon two point function in Fig. 1 (a) is expressed as

$$\Pi_{\mu\nu}^{AB}(p) = -N_f \int \frac{d^D k}{(2\pi)^D} \text{Tr} \left[(-g) i\gamma_{\mu} T_{ab}^A \frac{1}{i(\not{p} + \not{k})} (-g) i\gamma_{\nu} T_{ba}^B \frac{1}{i\not{k}} \right]. \quad (29)$$

After some algebra we obtain

$$\Pi_{\mu\nu}^{AB}(p) = -N_f \frac{g_0^2}{(4\pi)^2} \delta_{AB} (\delta_{\mu\nu} p^2 - p_{\mu} p_{\nu}) \left\{ \frac{2}{3} \frac{1}{\epsilon} + \frac{2}{3} \log(\mu^2/p^2) + \Delta_{\text{cont}} \right\} \quad (30)$$

with

$$\Delta_{\text{cont}} = \frac{10}{9} \quad (31)$$

where $1/\bar{\epsilon} = 1/\epsilon - \gamma + \ln(4\pi)$ and μ is the renormalization scale. In the $\overline{\text{MS}}$ scheme, the pole term $1/\bar{\epsilon}$ should be eliminated.

Let us turn to the calculation of the one loop corrections to the gluon self energy on the lattice. We illustrate the relevant diagrams in Fig. 1: (a), (b) for the contribution of the domain wall fermion and (c), (d) for the Pauli-Villars regulators. Each loop diagram is expressed by

$$\Pi_{\mu\nu}^{(a)AB}(p) = -N_f \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \text{Tr} \left[V_{1\mu}^A(k, p+k)_{uv} S_F(p+k)_{vs} V_{1\nu}^B(p+k, k)_{st} S_F(k)_{tu} \right], \quad (32)$$

$$\Pi_{\mu\nu}^{(b)AB}(p) = -N_f \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \text{Tr} \left[V_{2\mu\nu}^{AB}(k, k)_{st} S_F(k)_{ts} \right], \quad (33)$$

$$\Pi_{\mu\nu}^{(c)AB}(p) = +N_f \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \text{Tr} \left[\bar{V}_{1\mu}^A(k, p+k)_{uv} \bar{S}_F(p+k)_{vs} \bar{V}_{1\nu}^B(p+k, k)_{st} \bar{S}_F(k)_{tu} \right], \quad (34)$$

$$\Pi_{\mu\nu}^{(d)AB}(p) = +N_f \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \text{Tr} \left[\bar{V}_{2\mu\nu}^{AB}(k, k)_{st} \bar{S}_F(k)_{ts} \right], \quad (35)$$

where $\bar{V}_{1\mu}^A$, $\bar{V}_{2\mu\nu}^A$ and $\bar{S}_F$ denote the one-, two-gluon vertices and the propagator for the Pauli-Villars regulators. The sum of these contributions is expressed by $\Pi_{\mu\nu}^{AB}(p)$:

$$\Pi_{\mu\nu}^{AB}(p) = \Pi_{\mu\nu}^{(a)AB}(p) + \Pi_{\mu\nu}^{(b)AB}(p) + \Pi_{\mu\nu}^{(c)AB}(p) + \Pi_{\mu\nu}^{(d)AB}(p) \quad (36)$$

$$= \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} I_{\mu\nu}^{AB}(k, p). \quad (37)$$

Since the function $\Pi_{\mu\nu}^{AB}(p)$ has an infrared divergence for $p^2 \rightarrow 0$, we extract it by employing an analytically integrable expression $\tilde{I}_{\mu\nu}^{AB}(p)$ which has the same infrared behavior as $I_{\mu\nu}^{AB}(p)$,

$$\Pi_{\mu\nu}^{AB}(p) = \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \tilde{I}_{\mu\nu}^{AB}(k, p) + \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \left\{ I_{\mu\nu}^{AB}(k, p) - \tilde{I}_{\mu\nu}^{AB}(k, p) \right\}, \quad (38)$$

where

$$\tilde{I}_{\mu\nu}^{AB}(k, p) = \theta(\Lambda^2 - k^2) \text{Tr} \left[(-g) i \gamma_\mu T_{ab}^A \frac{1}{i(\not{p} + \not{k})} (-g) i \gamma_\nu T_{ba}^B \frac{1}{i\not{k}} \right], \quad (39)$$

with a cut-off Λ ($\leq \pi$). Now we find that the second term in the right-hand side of eq.(38) is regular in terms of p , we can make a Taylor expansion,

$$\Pi_{\mu\nu}^{AB}(p) = \int_{-\pi}^{\pi} \frac{d^4k}{(2\pi)^4} \tilde{I}_{\mu\nu}^{AB}(k, p)$$

$$\begin{aligned}
& + \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \left\{ I_{\mu\nu}^{AB}(k, 0) - \tilde{I}_{\mu\nu}^{AB}(k, 0) \right\} \\
& + \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} p_{\rho} \frac{\partial}{\partial p_{\rho}} \left\{ I_{\mu\nu}^{AB}(k, p) - \tilde{I}_{\mu\nu}^{AB}(k, p) \right\} \Big|_{p=0} \\
& + \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \frac{1}{2} p_{\rho} p_{\lambda} \frac{\partial^2}{\partial p_{\rho} \partial p_{\lambda}} \left\{ I_{\mu\nu}^{AB}(k, p) - \tilde{I}_{\mu\nu}^{AB}(k, p) \right\} \Big|_{p=0}, \tag{40}
\end{aligned}$$

where $\rho, \lambda = 1, 2, 3, 4$.

Let us consider each term in eq.(40). We first point out that the linear term in p is not allowed from the Lorentz symmetry. The integration of $\tilde{I}_{\mu\nu}^{AB}(k, p) - \tilde{I}_{\mu\nu}^{AB}(k, 0)$ is easily performed analytically,

$$\begin{aligned}
& \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \left\{ \tilde{I}_{\mu\nu}^{AB}(k, p) - \tilde{I}_{\mu\nu}^{AB}(k, 0) \right\} \\
& = -N_f \frac{g_0^2}{(4\pi)^2} \delta_{AB} \left[(\delta_{\mu\nu} p^2 - p_{\mu} p_{\nu}) \left\{ \frac{2}{3} \log(\Lambda^2/p^2) + \frac{5}{9} \right\} + \delta_{\mu\nu} p^2/3 \right]. \tag{41}
\end{aligned}$$

As for $\Pi_{\mu\nu}^{AB}(0) = \int_{-\pi}^{\pi} I_{\mu\nu}^{AB}(k, 0)$, we find that this function vanishes. This can be shown as follows. For the fermion contribution,

$$\Pi_{\mu\nu}^{(a)AB}(0) = +N_f \frac{g_0^2}{2} \delta_{AB} \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \text{Tr} [v_{1\mu}(k, k) S_F(k)_{ts} v_{1\nu}(k, k) S_F(k)_{st}], \tag{42}$$

$$= -N_f \frac{g_0^2}{2} \delta_{AB} \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \text{Tr} \left[v_{1\mu}(k, k) \frac{\partial}{\partial k_{\nu}} S_F(k)_{tt} \right], \tag{43}$$

$$= +N_f \frac{g_0^2}{2} \delta_{AB} \int_{-\pi}^{\pi} \frac{d^4 k}{(2\pi)^4} \text{Tr} \left[\frac{\partial}{\partial k_{\nu}} v_{1\mu}(k, k) S_F(k)_{tt} \right], \tag{44}$$

$$= -\Pi_{\mu\nu}^{(b)AB}(0), \tag{45}$$

where we use the identities

$$\frac{\partial}{\partial k_{\nu}} S_F^{-1}(k)_{st} = v_{1\nu}(k, k) \delta_{st}, \tag{46}$$

$$\frac{\partial}{\partial k_{\nu}} [S_F^{-1}(k)_{st} S_F(k)_{tu}] = 0, \tag{47}$$

$$\frac{\partial}{\partial k_{\nu}} v_{1\mu}(k, k) = v_{2\mu\nu}(k, k). \tag{48}$$

In a similar way we can show

$$\Pi_{\mu\nu}^{(c)AB}(0) = -\Pi_{\mu\nu}^{(d)AB}(0) \tag{49}$$

for the contribution of the Pauli-Villars regulator. After all, the sum of all the contributions vanishes:

$$\Pi_{\mu\nu}^{AB}(0) = \Pi_{\mu\nu}^{(a)AB}(0) + \Pi_{\mu\nu}^{(b)AB}(0) + \Pi_{\mu\nu}^{(c)AB}(0) + \Pi_{\mu\nu}^{(d)AB}(0) = 0. \tag{50}$$

We now turn to the last term in eq. (40). The calculation of this term requires tedious algebra and yields quite lengthy results that make it practically difficult to give their full expression here. However, we think that it would be instructive to explain why the Pauli-Villars regulator is required by showing a part of the calculation. Let us take the term proportional to $\text{Tr}[\gamma_\mu \gamma_\alpha \gamma_\nu \gamma_\beta]$ as an example,

$$I_{\mu\nu}^{(4\gamma)AB}(k, p) = -N_f g_0^2 \frac{\delta_{AB}}{2} \text{Tr}[\gamma_\mu \gamma_\alpha \gamma_\nu \gamma_\beta] \cos(k_\mu + p_\mu/2) \sin(k_\alpha + p_\alpha) \cos(k_\nu + p_\nu/2) \sin(k_\beta) \\ \times \left\{ G'_R(s, t) G_R(t, s) - \overline{G}'_R(s, t) \overline{G}_R(t, s) \right\}, \quad (51)$$

where G'_R and $\overline{G}'_R$ are functions of $p+k$, while G_R and $\overline{G}_R$ are functions of k . $\overline{G}_R$ is obtained from eq.(24) with $m = -1$. The fermion contribution is evaluated as

$$G'_R(s, t) G_R(t, s) = \sum_{s, t} \left[A' e^{-\alpha'|s-t|} + A'_L e^{\alpha'(s+t-2N_s)} + A'_R e^{\alpha'(-s-t+2)} \right] \\ \times \left[A e^{-\alpha|s-t|} + A_L e^{\alpha(s+t-2N_s)} + A_R e^{\alpha(-s-t+2)} \right] \quad (52) \\ = A' A \left\{ \frac{-2e^{-(\alpha'+\alpha)}}{[1 - e^{-(\alpha'+\alpha)}]^2} + N_s \frac{1 + e^{-(\alpha'+\alpha)}}{1 - e^{-(\alpha'+\alpha)}} \right\} \\ + A'(A_L + A_R) \frac{1 + e^{(\alpha'+\alpha)}}{1 - e^{(\alpha'+\alpha)}} \frac{e^{2\alpha}}{1 - e^{2\alpha}} \\ + (A'_L + A'_R) A \frac{1 + e^{(\alpha'+\alpha)}}{1 - e^{(\alpha'+\alpha)}} \frac{e^{2\alpha'}}{1 - e^{2\alpha'}} \\ + (A'_L A_L + A'_R A_R) \frac{e^{2(\alpha'+\alpha)}}{[1 - e^{(\alpha'+\alpha)}]^2} \quad (53)$$

with

$$A_L = -\frac{1 - W e^{-\alpha}}{1 - W e^{\alpha}} A, \quad (54)$$

$$A_R = -e^{-2\alpha} A, \quad (55)$$

where the superscript $\prime$ denotes the function of $p+k$. On the other hand, the Pauli-Villars part is given by

$$\overline{G}'_R(s, t) \overline{G}_R(t, s) = \sum_{s, t} A' \left[e^{-\alpha'|s-t|} - e^{\alpha'(s-t-N_s)} - e^{\alpha'(-s+t-N_s)} \right] \\ \times A \left[e^{-\alpha|s-t|} - e^{\alpha(s-t-N_s)} - e^{\alpha(-s+t-N_s)} \right] \quad (56)$$

$$= A' A \left\{ \frac{-2e^{-(\alpha'+\alpha)}}{[1 - e^{-(\alpha'+\alpha)}]^2} + N_s \frac{1 + e^{-(\alpha'+\alpha)}}{1 - e^{-(\alpha'+\alpha)}} + \frac{2e^{-(\alpha'+\alpha)}}{[1 - e^{-(\alpha'+\alpha)}]^2} \right\}, \quad (57)$$

where we find the exactly same $O(N_s)$ term as in the domain-wall fermion part. This is sustained even after taking the derivative in terms of p according to eq.(40). Through this

kind of analytical calculation we have checked that all the $O(N_s)$ terms of the fermion contribution in the last term of eq. (40) are canceled out by the Pauli-Villars one, which is proportional to N_s after collecting all the terms.

After all, eq. (40) is summarized as

$$\begin{aligned} \Pi_{\mu\nu}^{AB}(p) = & -N_f \frac{g_0^2}{(4\pi)^2} \delta_{AB} \left[(\delta_{\mu\nu} p^2 - p_\mu p_\nu) \left\{ \frac{2}{3} \log \left(\frac{\pi^2}{p^2 a^2} \right) + \frac{5}{9} \right\} + \delta_{\mu\nu} p^2 / 3 \right. \\ & \left. + A \delta_{\mu\nu} p^2 + B p_\mu p_\nu \right]. \end{aligned} \quad (58)$$

The momentum integration is carried out numerically employing the Monte Carlo routine BASES[22]. We choose $\Lambda = \pi$ for the cut-off. The results are checked by a mode sum for a periodic box of a size L^4 with $L = 128$ after transforming the momentum variable through $k'_\mu = k_\mu - \text{sink}_\mu$. The estimated values for $A + 1/3$ and B are presented in Table II as a function of M . We observe that the expected relation $A + 1/3 = -B$ is well satisfied within error bars. Finally we obtain

$$\Pi_{\mu\nu}^{AB}(p) = -N_f \frac{g_0^2}{(4\pi)^2} \delta_{AB} (\delta_{\mu\nu} p^2 - p_\mu p_\nu) \left[-\frac{2}{3} \log(p^2 a^2) + \Delta_{\text{latt}} \right], \quad (59)$$

where

$$\Delta_{\text{latt}} = A + \frac{8}{9} + \frac{2}{3} \log(\pi^2). \quad (60)$$

Comparing the above expression to the continuum counterpart in eq.(30), we obtain the finite renormalization constant d_f in eq.(1),

$$d_f = \frac{1}{16\pi^2} (\Delta_{\text{latt}} - \Delta_{\text{cont}}), \quad (61)$$

where the numerical values are given in Table II as a function of M . We find that the sign is opposite to and the magnitude is compatible with the Wilson, the clover and the Kogut-Susskind cases : $d_f = 0.0066949$ [23, 24], $d_f = 0.0314917$ [25], and $d_f = 0.0026248$ [26], respectively.

IV. THE RATIO OF CONTINUUM AND LATTICE Λ PARAMETERS

We first show how to obtain the mean-field improved $\overline{\text{MS}}$ coupling $g_{\overline{\text{MS}}}^2(\mu)$ at the scale μ from the lattice bare coupling g_0^2 . A detailed description for the quenched case is already given in Sec. VI of Ref. [14]. For the full QCD case the mean field improvement modifies

the relation of $g_{\overline{\text{MS}}}^2(\mu)$ and g_0^2 in eq.(1):

$$\frac{1}{g_{\overline{\text{MS}}}^2(\mu)} = \frac{P}{g_0^2} + d_g + c_p + \frac{22}{16\pi^2} \log(\mu a) + N_f \left(d_f - \frac{4}{48\pi^2} \log(\mu a) \right). \quad (62)$$

One may use an alternative formula[27]

$$\begin{aligned} \frac{1}{g_{\overline{\text{MS}}}^2(\mu)} = & \frac{c_0 P + 8c_1 R1 + 16c_2 R2 + 8c_3 R3}{g_0^2} \\ & + d_g + (c_0 \cdot c_p + 8c_1 \cdot c_{R1} + 16c_2 \cdot c_{R2} + 8c_3 \cdot c_{R3}) + \frac{22}{16\pi^2} \log(\mu a) \\ & + N_f \left(d_f - \frac{4}{48\pi^2} \log(\mu a) \right), \end{aligned} \quad (63)$$

where

$$P = \frac{1}{3} \text{Tr} U_{\text{plaquette}} = 1 - c_p g_0^2 + O(g_0^4), \quad (64)$$

$$R1 = \frac{1}{3} \text{Tr} U_{\text{rectangle}} = 1 - c_{R1} g_0^2 + O(g_0^4), \quad (65)$$

$$R2 = \frac{1}{3} \text{Tr} U_{\text{chair}} = 1 - c_{R2} g_0^2 + O(g_0^4), \quad (66)$$

$$R3 = \frac{1}{3} \text{Tr} U_{\text{parallelogram}} = 1 - c_{R3} g_0^2 + O(g_0^4), \quad (67)$$

and the measured values may be employed for P , $R1$, $R2$ and $R3$. Table I summarizes the values of c_p , c_{R1} , c_{R2} and c_{R3} for various gauge actions, which are taken from Table XVI of Ref. [14]. One may also use the mean-field estimate for M , which is given by $\tilde{M} = M - 4(1 - P^{1/4})$ [14]. No other modifications are necessary in this case: Just use d_f at $\tilde{M}$.

As discussed in Sec. I, we can determine the ratio of Λ_L and $\Lambda_{\overline{\text{MS}}}$ from the finite renormalization of the coupling constant. From eq.(7), the ratio is written as

$$\frac{\Lambda_{\overline{\text{MS}}}}{\Lambda_L} = \exp \left(-\frac{d_g}{2\beta_0} \right) \times \exp \left(-\frac{N_f d_f}{2\beta_0} \right), \quad (68)$$

where the first factor is given in Table I for the various gauge actions, and the second one is given in Table II as a function of M , in the case that $N_f = 0, 1, 2, 3, 4$.

V. CONCLUSION

In this paper we evaluate the fermion contribution to the finite renormalization of the coupling constant in the domain-wall QCD, which determines the ratio of the lattice Λ parameter to the continuum one. The numerical values are given as a function of M . We

show that the massive fermion modes emerging from the fermion loop contribution to the gluon self energy can be subtracted by the Pauli-Villars regulator. We explain how to implement the mean field improvement for the renormalized coupling $g_{\overline{\text{MS}}}^2(\mu)$ and the Λ parameter ratio.

Acknowledgments

This work is supported in part by the Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology. (Nos. 13135204, 14046202, 15204015, 15540251, 15740165).

-
- [1] Y. Shamir, Nucl. Phys. **B406** (1993) 90.
 - [2] V. Furman and Y. Shamir, Nucl. Phys. **B439** (1995) 54.
 - [3] RBC Collaboration, T. Izubuchi *et al.*, hep-lat/0210011
 - [4] A. Hasenfratz and P. Hasenfratz, Phys. Lett. **93B** (1980) 165; Nucl. Phys. **B193** (1981) 210.
 - [5] P. Weisz, Nucl. Phys. **B212** (1983) 1; P. Weisz and R. Wohlert, Nucl. Phys. **B236** (1984) 397; erratum, *ibid.* **B347** (1984) 544.
 - [6] Y. Iwasaki and S. Sakai, Nucl. Phys. **B248** (1984) 441.
 - [7] Y. Iwasaki and T. Yoshié, Phys. Lett. **143B** (1984) 449.
 - [8] S. Sakai, T. Saito and A. Nakamura, Nucl. Phys. **B584** (2000) 528.
 - [9] S. Aoki and Y. Taniguchi, Phys. Rev. **D59** (1999) 054510.
 - [10] S. Aoki and Y. Taniguchi, Phys. Rev. **D59** (1999) 094506.
 - [11] S. Aoki, T. Izubuchi, Y. Kuramashi and Y. Taniguchi, Phys. Rev. **D59** (1999) 094505.
 - [12] S. Aoki, T. Izubuchi, Y. Kuramashi and Y. Taniguchi, Phys. Rev. **D60** (1999) 114504.
 - [13] S. Aoki and Y. Kuramashi, Phys. Rev. **D63** (2001) 054504.
 - [14] S. Aoki, T. Izubuchi, Y. Kuramashi and Y. Taniguchi, Phys. Rev. **D67** (2003) 094502.
 - [15] P. Vranas, Phys. Rev. **D57** (1998) 1415.
 - [16] M. Lüscher and P. Weisz, Commun. Math. Phys. **97** (1985) 59; erratum, *ibid.* **98** (1985) 433.
 - [17] Y. Iwasaki, preprint, UTHEP-118 (Dec. 1983), unpublished.

- [18] K. G. Wilson, in *Recent Development of Gauge Theories*, eds. G. 'tHooft *et al.* (Plenum, New York, 1980).
- [19] T. Takaishi, Phys. Rev. **D54** (1996) 1050; P. de Forcrand *et al.*, Nucl. Phys. **B577** (2000) 263.
- [20] CP-PACS Collaboration, A. Ali Khan *et al.*, Phys. Rev. **D63** (2001) 114504; Phys. Rev. **D64** (2001) 114506; CP-PACS Collaboration, J. I. Noaki *et al.*, hep-lat/0108013.
- [21] RBC Collaboration, Y. Aoki *et al.*, Nucl. Phys. Proc. Suppl. **106** (2002) 245; RBC Collaboration, K. Orginos *et al.*, Nucl. Phys. Proc. Suppl. **106** (2002) 721; Y. Aoki *et al.*, hep-lat/0211023.
- [22] S. Kawabata, Comput. Phys. Commun. **41** (1986) 127; **88** (1995) 309.
- [23] P. Weisz, Phys. Lett. **100B** (1981) 331.
- [24] H. Kawai, R. Nakayama and K. Seo, Nucl. Phys. **B189** (1981) 40.
- [25] S. Sint and R. Sommer, Nucl. Phys. **B465** (1996) 71.
- [26] H. S. Sharatchandra, H. J. Thun and P. Weisz, Nucl. Phys. **B192** (1981) 205.
- [27] A. Ali Khan *et al.*, Phys. Rev. **D65** (2002) 054505.

TABLE I: Values of the perturbative quantities d_g , c_P , and c_{Ri} for the various gauge actions. Gluon contribution to the Λ parameter ratio in eq.(68) is given for the case of $N_f = 0, 1, 2, 3, 4$.

action	c_1	c_3						$\exp(-d_g/(2\beta_0))$				
			d_g	c_p	c_{R1}	c_{R2}	c_{R3}	$N_f = 0$	$N_f = 1$	$N_f = 2$	$N_f = 3$	$N_f = 4$
Plaquette	0	0	-0.4682	1/3	0.5747	0.5228	0.5687	28.81(2)	35.78(3)	45.80(4)	60.80(5)	84.45(8)
Iwasaki	-0.331	0	0.1053	0.1401	0.2689	0.2223	0.2465	0.4696(3)	0.4473(3)	0.4231(3)	0.3970(3)	0.3687(3)
DBW2	-1.40686	0	0.5317	0.0511	0.1040	0.0818	0.0921	0.02200(2)	0.01720(1)	0.01300(1)	0.009423(8)	0.006488(6)
Symanzik	-1/12	0	-0.2361	0.2440	0.4417	0.3846	0.4215	5.445(4)	6.074(5)	6.879(6)	7.935(7)	9.365(9)
Iwasaki'	-0.27	-0.04		0.1471	0.2797	0.2342	0.2611					
Wilson	-0.252	-0.17	0.1196	0.1286	0.2439	0.2070	0.2352	0.4238(3)	0.4010(3)	0.3765(3)	0.3502(3)	0.3220(3)

TABLE II: Numerical values for $A + 1/3$, B and d_f as a function of M Fermion contribution to the Λ parameter ratio in eq.(68) is given for the case of $N_f = 1, 2, 3, 4$.

M	$A + 1/3$	B	d_f	$\exp(-N_f d_f / (2\beta_0))$			
				$N_f = 1$	$N_f = 2$	$N_f = 3$	$N_f = 4$
0.05	-5.4253(27)	5.4259(27)	-0.028209(17)	1.24054(16)	1.58537(44)	2.10102(95)	2.9127(19)
0.10	-4.6075(22)	4.6092(23)	-0.023030(14)	1.19240(13)	1.45676(33)	1.83331(68)	2.3937(13)
0.15	-4.1408(20)	4.1414(19)	-0.020075(13)	1.16578(11)	1.38810(29)	1.69612(57)	2.1400(10)
0.20	-3.8102(18)	3.8106(18)	-0.017981(11)	1.14728(10)	1.34143(25)	1.60520(48)	1.97679(85)
0.25	-3.5497(18)	3.5494(18)	-0.016332(11)	1.132911(97)	1.30577(24)	1.53700(45)	1.85699(79)
0.30	-3.3341(15)	3.3346(16)	-0.0149658(97)	1.121148(83)	1.27696(20)	1.48273(38)	1.76331(65)
0.35	-3.1475(15)	3.1483(14)	-0.0137846(96)	1.111075(81)	1.25255(20)	1.43734(36)	1.68612(61)
0.40	-2.9819(14)	2.9825(13)	-0.0127360(87)	1.102208(73)	1.23128(17)	1.39822(32)	1.62042(53)
0.45	-2.8310(13)	2.8316(12)	-0.0117805(83)	1.094190(69)	1.21221(16)	1.36349(30)	1.56279(49)
0.50	-2.6925(12)	2.6938(12)	-0.0109029(79)	1.086878(65)	1.19496(15)	1.33236(28)	1.51167(45)
0.55	-2.5635(12)	2.5650(12)	-0.0100864(76)	1.080117(62)	1.17912(15)	1.30403(26)	1.46560(42)
0.60	-2.4418(12)	2.4433(11)	-0.0093153(73)	1.073772(60)	1.16436(14)	1.27784(25)	1.42339(40)
0.65	-2.3272(11)	2.3282(11)	-0.0085901(67)	1.067839(55)	1.15064(13)	1.25368(22)	1.38480(35)
0.70	-2.2176(10)	2.2182(11)	-0.0078955(64)	1.062186(52)	1.13767(12)	1.23097(21)	1.34883(33)
0.75	-2.11237(99)	2.11208(98)	-0.0072294(63)	1.056794(51)	1.12535(12)	1.20958(20)	1.31520(31)
0.80	-2.01129(98)	2.01132(97)	-0.0065893(62)	1.051638(50)	1.11365(11)	1.18937(19)	1.28368(30)
0.85	-1.91395(88)	1.91450(89)	-0.0059729(55)	1.046696(44)	1.10249(10)	1.17023(17)	1.25404(26)
0.90	-1.82014(87)	1.82057(91)	-0.0053788(55)	1.041956(44)	1.091844(98)	1.15207(17)	1.22612(26)
0.95	-1.73060(81)	1.73063(86)	-0.0048118(51)	1.037451(40)	1.081777(90)	1.13501(15)	1.20005(23)
1.00	-1.64359(81)	1.64365(82)	-0.0042608(51)	1.033092(41)	1.072083(90)	1.11867(15)	1.17525(23)
1.05	-1.56012(77)	1.55976(76)	-0.0037322(49)	1.028928(39)	1.062866(85)	1.10321(14)	1.15194(21)
1.10	-1.48027(71)	1.47976(71)	-0.0032266(45)	1.024961(35)	1.054123(78)	1.08863(13)	1.13008(19)
1.15	-1.40379(68)	1.40313(69)	-0.0027422(43)	1.021174(33)	1.045815(73)	1.07484(12)	1.10952(18)
1.20	-1.33232(65)	1.33289(66)	-0.0022896(41)	1.017649(32)	1.038112(70)	1.06211(12)	1.09065(17)
1.25	-1.26633(62)	1.26646(62)	-0.0018718(39)	1.014405(30)	1.031050(66)	1.05050(11)	1.07352(16)
1.30	-1.20519(59)	1.20600(60)	-0.0014846(37)	1.011408(29)	1.024549(63)	1.03985(10)	1.05788(15)
1.35	-1.15142(57)	1.15256(57)	-0.0011441(36)	1.008780(28)	1.018865(60)	1.030569(98)	1.04431(14)
1.40	-1.10734(55)	1.10903(55)	-0.0008650(35)	1.006631(27)	1.014230(58)	1.023026(94)	1.03332(14)
1.45	-1.07434(53)	1.07514(54)	-0.0006560(34)	1.005025(26)	1.010774(56)	1.017416(90)	1.02517(13)
1.50	-1.05539(52)	1.05533(52)	-0.0005360(33)	1.004104(25)	1.008794(55)	1.014206(89)	1.02052(13)
1.55	-1.05344(53)	1.05437(53)	-0.0005236(33)	1.004009(26)	1.008590(55)	1.013876(89)	1.02004(13)
1.60	-1.07444(54)	1.07497(54)	-0.0006566(34)	1.005030(26)	1.010784(56)	1.017431(91)	1.02520(13)
1.65	-1.12617(56)	1.12665(56)	-0.0009842(36)	1.007548(27)	1.016207(59)	1.026241(96)	1.03800(14)
1.70	-1.21873(61)	1.21856(61)	-0.0015704(39)	1.012071(30)	1.025985(65)	1.04220(11)	1.06132(16)
1.75	-1.36869(68)	1.36952(68)	-0.0025200(43)	1.019442(34)	1.042025(74)	1.06857(12)	1.10022(18)
1.80	-1.60462(80)	1.60601(80)	-0.0040140(51)	1.031146(40)	1.067771(89)	1.11143(15)	1.16431(22)
1.85	-1.97527(99)	1.97602(99)	-0.0063612(63)	1.049807(50)	1.10951(11)	1.18225(19)	1.27263(30)
1.90	-2.5818(13)	2.5816(13)	-0.0102019(82)	1.081071(68)	1.18135(16)	1.30800(28)	1.47203(46)
1.95	-3.7162(19)	3.7153(19)	-0.017386(12)	1.14207(10)	1.32845(26)	1.58024(49)	1.93268(86)




FIG. 1: One loop corrections to the gluon self energy. Solid line is for the propagator of the domain-wall fermion and broken one for the Pauli-Villars regulator.